

“Assisted cloning” and “orthogonal-complementing” of an unknown state

Arun Kumar Pati

SEECs, Dean Street, University of Wales, Bangor LL 57 1UT, UK

(February 9, 2008)

We propose a protocol where one can exploit dual quantum and classical channels to achieve perfect “cloning” and “orthogonal-complementing” of an unknown state with a minimal assistance from a state preparer (without revealing what the input state is). The first stage of the protocol requires usual teleportation and in the second stage, the preparer disentangles the left over entangled states by a single particle measurement process and communicates a number of classical bits (1-cbit per copy) to different parties so that perfect copies and complement copies are produced. We discuss our protocol for producing two copies and three copies (and complement copies) using two and four particle entangled state and suggest how to generalise this for N copies and complement copies using multiparticle entangled state.

PACS NO: 03.67.Hk, 03.67.-a, 03.65.Bz

email:akpati@sees.bangor.ac.uk

Manipulation and extraction of quantum information are important task in the ongoing field of information theory. One of the interesting question raised by Wootters and Zurek [1] and Dieks [2] is whether it is possible to copy a quantum state perfectly. It was found that linearity of quantum theory does not allow us to do so. In recent years this question has again become important. Especially, in the field of quantum communication and quantum computing one needs to know the ability to process the quantum information content of a system. In the literature one finds another proof of a “no-cloning theorem” which is based only on unitarity of the evolution. This says that deterministic cloning of non-orthogonal states is impossible [3–5]. Even though we can not perfectly copy an unknown state, nevertheless, the possibility of producing approximate copies of a quantum state has been considered by Bužek and Hillery [6]. They [7] have shown that if two non-orthogonal states are cloned there will be errors introduced in the copies. Bužek *et al* [8] have discussed a network for quantum copying where one can realise a Universal Quantum Copying Machine (UQCM). Bruß *et al* [9] have constructed an optimal state dependent cloning machine for producing two clones. There has been a proposal by Duan and Guo [10] for probabilistic cloning of non-orthogonal quantum state using unitary operator and measurement process. The possibility of producing M copies from a given N input states has been investigated by Gisin and Masar [11]. The possibility of obtaining asymmetric clones were investigated by Cerf [12]. The problem of state discrimination and exact cloning has been discussed by Chefles and Barnett [13]. They have also discussed a network which interpolates between deterministic (approximate) and probabilistic (exact) cloning [14] from a given M initial states. Recently, we have asked a question if there exists a cloning machine, which can produce a linear superposition of multiple clones and found that such an ideal machine can not exist [15]. However, we have proved that if the machine fails some times it is possible to build a novel cloning machine, which can produce linear superposition of multiple clones [16]. We have also argued that the deterministic and probabilistic cloning machines are special cases of our novel cloning machine. This would find potential application in quantum information processing as it would provide a parallel quantum operation over many clones.

Recently, another limitation to quantum information processing has been found. This says that it is not possible to produce a conjugate copy of an unknown state using either linearity or unitarity of the quantum process [17]. The process of orthogonal-complementing of an unknown state is creating a state which is orthogonal to the unknown state. Independently, it is found that it is not possible to perfectly create an orthogonal-complement state of an unknown state, because the operation involves anti-unitary operator which can not be implemented in the laboratory on an isolated system [18]. In the case of a qubit a complement state can be related to a conjugate state by a rotation operator. Hence, these problems are identical (upto a rotation operator). The optimal manipulation of an unknown state with a universal NOT gate has been investigated in [18]. A related question on spin flipping has been investigated by Gisin and Popescu [19] who found that it is not possible to flip an arbitrary spin.

Although, there have been immense theoretical ideas about how well one can copy a quantum state, none of them seems to yield a perfect copy of the input state. Here, we should mention that in the Duan-Guo [10] type of probabilistic cloning machine one can produce perfect copies with some probabilities. The probabilities of success

obey certain inequalities which depend on the inner product of the input states. However, in the present scheme the perfect copies are produced with a probability independent of the input state. Moreover, there is no proof yet that using unitary and reduction operations one can produce an orthogonal complement copy of an unknown state. Also, there is no hope that one can test all the existing theoretical ideas experimentally where one can create a copy with some error.

The purpose of this paper is to investigate the possibility of copying and complementing an unknown state perfectly using resources like entangled states, Bell-state measurement, single particle von Neumann measurement and classical information. We go beyond the traditional cloning ideas that exist in the literature. The question we raise is this: Can we produce a perfect copy of an unknown quantum state with a minimal assistance from the state preparer (call him Victor). It turns out that with the help of Victor our protocol (which allows unitary and non-unitary (measurements) operations) can produce a perfect copy and a complement copy. In a strict sense a quantum cloning machine is not supposed to communicate with Victor because one would say that since Victor knows what the state is he could create as many copies as he wants and send them to the interested parties. Alternately, Victor could send his recipe to some-one to prepare the state which requires double infinity of bits of information [20] to be sent across a classical channel. But if for some reason Victor does not want to do that what can we do at best? To be more practical, we pose the question: If Victor has only one input state and does not want to send infinite of bits of information, then without violating any principles of quantum theory and *using quantum mechanical operations can we go around “no-cloning theorem”*? If Victor agrees to assist the parties with a minimum help without revealing what the input state is, then what we can do is described in this paper. It is shown that instead of sending infinite bits of information Victor can use the entangled state left after the teleportation process [21] and send one classical bit to Alice to create either a copy or an orthogonal-complement copy of the unknown state. Like in the teleportation process one qubit can be passed by sending two cbits and the remaining flows across the entanglement channel, similarly here (in the second stage of protocol), the infinite amount of bits of information can be passed to a distant site by just sending one cbit and remaining bits flow across the entanglement channel. This is a non-trivial observation in this context which must be remembered and should not be confused with the issue of what Victor could do in principle. (What a state preparer could do in principle is not our concern here, what he is doing in the present protocol is important.) The importance of this approach can be in those situations, where Victor is cut-off (one way) from the rest of the world and he can not send any quantum states but has a classical channel to send only cbits. An example could be, suppose Victor had owned a private company (“qubit company”) which produces qubits and was sending to interested parties. One day, his company is crashed and he has no resources to start his company again nor has he capitals. Then, if the other interested parties can send him one half of the particles from an entangled source, then they can be benefited from getting a copy or a complement copy of an unknown state after receiving 1 cbit from Victor. This way many parties can benefit from Victor, although he has lost his “qubit company”. But his knowledge and communication of 1 cbit per copy is quite helpful in such scenarios. The machine which produces a perfect copy and a complement copy of an unknown state with the minimal assistance, we call “assisted-cloning *cum* complementing machine” (ACCM). Here, by “minimal assistance” we mean any classical communication from Victor to Alice or Bob *without telling what the unknown state is*.

Suppose we have a pure input state $|\Psi\rangle_1 \in \mathcal{H} = C^2$, which is a qubit. An arbitrary qubit can be represented as

$$|\Psi\rangle_1 = \alpha|0\rangle_1 + \beta|1\rangle_1, \quad (1)$$

where we can choose α to be real and β to be a complex number, in general. This qubit can be represented by a point on a sphere S^2 (which is the projective Hilbert space for any two-state system) with the help of two real parameters θ and ϕ , where $\alpha = \cos \frac{\theta}{2}$ and $\beta = \sin \frac{\theta}{2} \exp(i\phi)$. Now Victor prepares the input state and sends it through a “ACCM” to produce two output states and later he can verify how good these copies are. Let Alice and Bob share one half of the particles from an EPR source as in the quantum teleportation protocol [21]. The EPR state for the particles 2 and 3 is given by,

$$|\Psi^-\rangle_{23} = \frac{1}{\sqrt{2}}(|01\rangle_{23} - |10\rangle_{23}). \quad (2)$$

Suppose Alice is in possession of 2 and Bob is in possession of 3. The input state $|\Psi\rangle_1$ is unknown to both Alice and Bob. The combined state of the unknown state 1 and the EPR state 23 (after reexpressing in terms of Bell-basis of particle 1 and 2) is

$$\begin{aligned} |\Psi\rangle_{123} &= |\Psi\rangle_1 \otimes |\Psi^-\rangle_{23} = -\frac{1}{2} [|\Psi^+\rangle_{12}(\sigma_z)|\Psi\rangle_3 \\ &+ |\Psi^-\rangle_{12}|\Psi\rangle_3 + |\Phi^+\rangle_{12}(i\sigma_y)|\Psi\rangle_3 + |\Phi^-\rangle_{12}(-\sigma_x)|\Psi\rangle_3], \end{aligned} \quad (3)$$

where σ 's are usual Pauli spin matrices generating $SU(2)$ Lie algebra. Once Alice performs a Bell-state measurement onto two-particle state 1 and 2, the resulting state is a tensor product of an entangled state between particle 1 and 2 and the outgoing state (teleported state up to a rotation operator). Suppose the measurement outcome of Alice is $|\Psi^-\rangle_{12}$, (the probability of this outcome is only $\frac{1}{4}$) then the resulting three particle state is given by

$$|\Psi^-\rangle_{12}\langle\Psi^-\Psi\rangle_{123} = -\frac{1}{2}|\Psi^-\rangle_{12} \otimes |\Psi\rangle_3. \quad (4)$$

When Alice communicates her measurement results to Bob (classical two bits of information) then Bob knows that he has received the original state. If the measurement outcome of Alice is other than $|\Psi^-\rangle_{12}$, then Bob has to apply suitable rotation operator to get the original state. Now can we create a copy of the original state at Alice's place? Since the teleportation process obeys the "no-cloning theorem", at first instance it might look impossible to have a copy at Alice's place. However, this impossibility becomes a possibility when we allow *one bit of classical information* from Victor to Alice.

Usually it is said that the Bell-state measurement destroys the particle 1 and 2, so there is no state available to Alice. This is some times stressed emphatically that the quantum teleportation process obeys "no-cloning" theorem, so particle 1 is destroyed. But careful analysis shows that obeying the "no-cloning theorem" of quantum theory and making two-particle detection (Bell-state measurements) are two different things. The point to be noted is that, in general the particles 1 and 2 need not be destroyed after Bell-state measurement. It so happens that when one uses photons for Bell-state analysis as was done by Bouwmeester et al [22] they are absorbed by the detectors and hence destroyed. For other particles this need not be so. Because the projection postulate says that when we apply a Bell-basis projector onto a the combined state $|\Psi\rangle_{123}$ indeed a Bell-state remains formally (see eqn.(4)). Quantum theory does not say that the particle 1 and 2 need to be destroyed after a measurement. How to implement this in practice is another question. A possible way to do this is through Bell-state analysis using quantum-nondemolition measurement (QND) [23] in the case of photons. This would need a device which can perform photon number QND measurement at single photon level. By sending particles 1 and 2 along with probe modes in a nonlinear Kerr medium one may think of a QND measurement. This would be, indeed, a challenge for experimentalist in the future. For other entangled sources such as spin-1/2 particles or two-level atoms one needs some suitable detector which can distinguish all four Bell-states and allow them to propagate freely for further processing. Let there be particles 1 and 2 after Bell-state measurement, which are in a singlet state. Alice sends particle 1 to Victor and keeps particle 2 in her possession. Now can Victor do some thing (without really telling what the input state is) so that Alice gets a copy of the unknown state? We show that this is possible in principle. Victor carries out another measurement on particle 1 by putting a linear polariser (in case of photon) or Stern-Gerlach apparatus (in case of spin 1/2 particles) to measure the state in another basis $\{|x\rangle, |y\rangle\}$, where the new basis is related to the old basis $\{|0\rangle, |1\rangle\}$ in the following manner

$$\begin{aligned} |0\rangle_1 &= \alpha|x\rangle_1 + \beta|y\rangle_1 \\ |1\rangle_1 &= \beta^*|x\rangle_1 - \alpha|y\rangle_1. \end{aligned} \quad (5)$$

Notice that the normalisation and orthogonality relation between basis vectors are preserved under this transformation. Interestingly, we find that the basis $|x\rangle_1 = |\Psi\rangle_1$ and the basis $|y\rangle_1 = |\Psi_\perp\rangle_1$, where $|\Psi_\perp\rangle_1 = (\alpha|1\rangle_1 - \beta^*|0\rangle_1)$ is the orthogonal-complement state to $|\Psi\rangle_1$. However, we keep $|x\rangle_1, |y\rangle_1$ for Victor just to distinguish the fact that he knows the state. When we write $|\Psi\rangle$ and $|\Psi_\perp\rangle$ for other particles we mean they are unknown to the parties concerned. Now writing the entangled state $|\Psi^-\rangle_{12}$ in the basis $|x\rangle_1, |y\rangle_1$ gives us

$$|\Psi^-\rangle_{12} = \frac{1}{\sqrt{2}}[|x\rangle_1(\alpha|1\rangle_2 - \beta^*|0\rangle_2) + |y\rangle_1(\alpha|0\rangle_2 + \beta|1\rangle_2)]. \quad (6)$$

If the outcome of Victor is $|y\rangle_1$, then he sends his measurement result (one bit of classical information) to Alice. Alice knows that her state of particle 2 has been found in the original state $(\alpha|0\rangle_2 + \beta|1\rangle_2)$ which is just a copy.

More explicitly, the total state after a Bell-state measurement and a single particle von-Neumann measurement is given by

$$|y\rangle_1\langle y|\Psi^-\rangle_{12}\langle\Psi^-\Psi\rangle_{123} = -\frac{1}{2\sqrt{2}}|y\rangle_1 \otimes |\Psi\rangle_2 \otimes |\Psi\rangle_3. \quad (7)$$

If the outcome of Victor's measurement result is $|x\rangle_1$ then the classical communication from Victor would tell Alice that she has obtained a state which is $(\alpha|1\rangle_2 - \beta^*|0\rangle_2)$. This is a complement copy of the unknown state. Now the state (if the outcome is $|x\rangle_1$) is given by

$$|x\rangle_1 \langle x| \Psi^-\rangle_{12} \langle \Psi^-| \Psi\rangle_{123} = -\frac{1}{2\sqrt{2}} |x\rangle_1 \otimes |\Psi_\perp\rangle_2 \otimes |\Psi\rangle_3. \quad (8)$$

On the other hand, if the measurement outcome of Alice is other than $|\Psi^-\rangle_{12}$ in the first stage of the protocol, then the result of the second stage of the protocol can be worked out in detail. First, we note that though the singlet state is same in any basis, the other Bell-states are not the same. In the basis $\{|x\rangle_1, |y\rangle_1\}$ we can express the other three Bell-states as

$$\begin{aligned} |\Psi^+\rangle_{12} &= -\frac{1}{\sqrt{2}} [|x\rangle_1 (\sigma_z) |\Psi_\perp\rangle_2 + |y\rangle_1 (\sigma_z) |\Psi\rangle_2] \\ |\Phi^+\rangle_{12} &= \frac{1}{\sqrt{2}} [|x\rangle_1 (i\sigma_y) |\Psi_\perp\rangle_2 + |y\rangle_1 (i\sigma_y) |\Psi\rangle_2] \\ |\Phi^-\rangle_{12} &= \frac{1}{\sqrt{2}} [|x\rangle_1 (\sigma_x) |\Psi_\perp\rangle_2 + |y\rangle_1 (\sigma_x) |\Psi\rangle_2] \end{aligned} \quad (9)$$

When Alice's outcome is $|\Psi^+\rangle_{12}$ in the first stage of the protocol, then the resulting states after a Bell-state measurement, a single particle von Neumann measurement and classical communication are given by

$$\begin{aligned} |x\rangle_1 \langle x| \Psi^+\rangle_{12} \langle \Psi^+| \Psi\rangle_{123} &= \frac{1}{2\sqrt{2}} |x\rangle_1 \otimes (\sigma_z) |\Psi_\perp\rangle_2 \otimes (\sigma_z) |\Psi\rangle_3 \\ |y\rangle_1 \langle y| \Psi^+\rangle_{12} \langle \Psi^+| \Psi\rangle_{123} &= \frac{1}{2\sqrt{2}} |y\rangle_1 \otimes (\sigma_z) |\Psi\rangle_2 \otimes (\sigma_z) |\Psi\rangle_3. \end{aligned} \quad (10)$$

Similarly, when Alice's outcome is $|\Phi^+\rangle_{12}$, then the resulting states after a Bell-state measurement, a single particle von Neumann measurement and classical communication are given by

$$\begin{aligned} |x\rangle_1 \langle x| \Phi^+\rangle_{12} \langle \Phi^+| \Psi\rangle_{123} &= -\frac{1}{2\sqrt{2}} |x\rangle_1 \otimes (i\sigma_y) |\Psi_\perp\rangle_2 \otimes (i\sigma_y) |\Psi\rangle_3 \\ |y\rangle_1 \langle y| \Phi^+\rangle_{12} \langle \Phi^+| \Psi\rangle_{123} &= -\frac{1}{2\sqrt{2}} |y\rangle_1 \otimes (i\sigma_y) |\Psi\rangle_2 \otimes (i\sigma_y) |\Psi\rangle_3. \end{aligned} \quad (11)$$

Finally, when Alice's outcome is $|\Phi^-\rangle_{12}$, then the resulting states after a Bell-state measurement, a single particle von Neumann measurement and classical communication are given by

$$\begin{aligned} |x\rangle_1 \langle x| \Phi^-\rangle_{12} \langle \Phi^-| \Psi\rangle_{123} &= \frac{1}{2\sqrt{2}} |x\rangle_1 \otimes (\sigma_x) |\Psi_\perp\rangle_2 \otimes (\sigma_x) |\Psi\rangle_3 \\ |y\rangle_1 \langle y| \Phi^-\rangle_{12} \langle \Phi^-| \Psi\rangle_{123} &= \frac{1}{2\sqrt{2}} |y\rangle_1 \otimes (\sigma_x) |\Psi\rangle_2 \otimes (\sigma_x) |\Psi\rangle_3. \end{aligned} \quad (12)$$

Thus, (7),(8) and (10-12) are exact ones and the main results of this paper. Our analysis illustrates the power of Bell-state and single particle measurements. The important observation is that if after the Bell-state measurement Victor finds $|y\rangle_1$ in a single particle measurement, then 1-cbit from Victor to Alice will result in an exact copy or a copy up to a rotation operator at Alice's place. If Victor finds $|x\rangle_1$, then sending of 1-cbit will yield an exact replica of complement state or a complement state up to a rotation operator. Interestingly, the rotation operators that Alice has to apply to get a copy is same as that of the Bob's case to get the original state.

In the special case, if we restrict our unknown state to be real, i.e., $|\Psi\rangle = \cos\theta|0\rangle + \sin\theta|1\rangle$, which means on the projective Hilbert space S^2 the point lies on the equatorial line, then the azimuthal angle is zero. Alice just has to perform a rotation or do nothing after receiving the classical information from Victor in case the measurement outcome is $|x\rangle_1$ or $|y\rangle_1$ and in both cases she gets a copy of the unknown state. Thus for a real unknown state our protocol produces a clone 100% of the time. For an arbitrary unknown state this protocol is able to produce an accurate copy of the original input state 50% of the time and an orthogonal-complement copy 50% of the time. Had our protocol worked only in the case of $|\Psi^-\rangle$, then the probability to produce a clone would have been $\frac{1}{8}$. But in the present case our protocol works for all Bell-state outcomes, hence probability to produce a copy is $\frac{1}{2}$, as mentioned. The fidelity of getting the copy and the complement copy is *maximum*, i.e. *unit*. One may tend to think that since the preparer is communicating with Alice this protocol may not qualify to be called quantum cloning. However, note that although Victor communicates the measurement result on some basis he is not going to tell what the exact value of α and β is. The measurement result sent by Victor does not contain all information about the unknown state. In

fact, the state of the particle 1 is no longer the original one, it is a member of the entangled pair. Knowledge of the unknown state is still not revealed to Alice and Bob. So this is not a classical cloning machine.

Can we imagine producing more copies or complement copies using the “assisted-quantum cloning” and “complementing” protocol? That is, given an input state $|\Psi\rangle$ can we have an output state $|\Psi\rangle \otimes |\Psi\rangle \otimes |\Psi\rangle$ or more? At first sight it may seem that may be if we use three particle entanglement we could generate *one to three* copies at different sites with the help of Victor. But it turns out that with a three-particle entangled source one can again produce only one perfect copy or one complement copy. So this analysis we are not going to present. The useful resource for producing *one to three* copies or 2 copies and a complement copy is a four-particle entangled state of the type Greenberger-Horne-Zeilinger (GHZ) [24]. This four particle state was originally proposed in the context of decay of four spin-half particles and was shown to yield non-local correlations without inequalities. The four particle entangled state (GHZ state) is given by

$$|\Psi\rangle_{2345} = \frac{1}{\sqrt{2}}(|0011\rangle_{2345} + |1100\rangle_{2345}) \quad (13)$$

Suppose we have three parties Alice, Bob and Carla who share the particles from this entangled state. Alice has particle 2, Bob has particles 3 and 4 and Carla has particle 5. Let Victor prepare a state which is unknown to Alice, Bob and Carla and is given by (1). He sends particle 1 to Alice who is now in possession of two particles 1 and 2. Now the combined five particle state is given by $|\Psi\rangle_1 \otimes |\Psi\rangle_{2345}$. Let us express the basis of states of particle 1 and 2 in their respective Bell-basis. Then the total state can be written as

$$\begin{aligned} |\Psi\rangle_{12345} &= |\Psi\rangle_1 \otimes |\Psi\rangle_{2345} \frac{1}{2} [|\Psi^+\rangle_{12}(\beta|011\rangle_{345} - \alpha|100\rangle_{345}) \\ &\quad - |\Psi^-\rangle_{12}(\beta|011\rangle_{345} + \alpha|100\rangle_{345}) + |\Phi^+\rangle_{12}(\alpha|011\rangle_{345} - \beta|100\rangle_{345}) \\ &\quad + |\Phi^-\rangle_{12}(\alpha|011\rangle_{345} + \beta|100\rangle_{345})] \end{aligned} \quad (14)$$

Now Alice carries out a Bell-state measurement on the particle 1 and 2. The two-particle projection would yield any one of four possible results. If the readout of Alice’s measurement is $|\Psi^-\rangle_{12}$ then the resulting state will be

$$|\Psi^-\rangle_{12} \langle \Psi^- | \Psi \rangle_{12345} = -\frac{1}{2} |\Psi^-\rangle_{12} (\beta|011\rangle_{345} + \alpha|100\rangle_{345}) \quad (15)$$

Thus the Bell-state measurement leaves particle 1 and 2 in a maximally entangled state and particle 3,4 and 5 in a three-particle entangled state. After the above measurement Alice sends her results via classical channel two bits of information to both Bob and Carla. In next step, Bob who is in possession of particle 3 and 4, carries out another Bell-state measurement on them. Let us express the above state in terms of Bell-states of 3 and 4. This is given by

$$\begin{aligned} |\Psi^-\rangle_{12} \langle \Psi^- | \Psi \rangle_{12345} &= -\frac{1}{2} |\Psi^-\rangle_{12} [|\Psi^+\rangle_{34}(\alpha|0\rangle_5 + \beta|1\rangle_5) \\ &\quad - |\Psi^-\rangle_{34}(\alpha|0\rangle_5 - \beta|1\rangle_5)] \end{aligned} \quad (16)$$

After a Bell-state measurement onto particle 3 and 4 if the readout is (say) $|\Psi^+\rangle_{34}$, then the state of particle 5 is found to be in the original state. If the read out is $|\Psi^-\rangle_{34}$ then the state of particle 5 is nothing but the original state up to a rotation operator σ_z . Suppose Bob’s measurement gives a result $|\Psi^+\rangle_{34}$, then the classical information from Bob to Carla would yield the state

$$|\Psi^+\rangle_{34} \langle \Psi^+ | \Psi^- \rangle_{12} \langle \Psi^- | \Psi \rangle_{12345} = -\frac{1}{2} |\Psi^-\rangle_{12} \otimes |\Psi^+\rangle_{34} \otimes |\Psi\rangle_5 \quad (17)$$

Note that in the second stage Bob needs to send only one bit of classical information to Carla as he could get only two possible Bell-state measurement results. The resulting state at Carla’s hand is nothing but the teleported state of the original input state (this can be called teleportation of an unknown state using four-particle entangled state, which has not been discussed in the literature). Recently, we find Karlsson and Bourennane [25] have discussed teleportation of an unknown state using three particle entangled state. But as we have said, three-particle entanglement is not a useful resource for cloning and complementing unknown states. After teleportation of the original state the particle 1 and 2 and 3 and 4 are in a maximally entangled state. Now Alice and Bob send particle 1 and 3 to Victor one after the other. When Victor gets the particles (1 and 3) he chooses to measure the states in the basis $|x\rangle_i, |y\rangle_i, (i = 1, 3)$, where

$$\begin{aligned}
|0\rangle_i &= \alpha|x\rangle_i + \beta|y\rangle_i \\
|1\rangle_i &= \beta^*|x\rangle_i - \alpha|y\rangle_i
\end{aligned} \tag{18}$$

In the new basis the total state is given by

$$\begin{aligned}
&|\Psi^+\rangle_{34}\langle\Psi^+|\Psi^-\rangle_{12}\langle\Psi^+|\Psi\rangle_{12345} = \\
&\frac{1}{4}[[x]_3(\alpha|1\rangle_4 + \beta^*|0\rangle_4) - |y\rangle_3(\alpha|0\rangle_4 - \beta|1\rangle_4)] \otimes \\
&[|x\rangle_1(\alpha|1\rangle_2 - \beta^*|0\rangle_2) + |y\rangle_1(\alpha|0\rangle_2 + \beta|1\rangle_2)] \otimes |\Psi\rangle_5
\end{aligned} \tag{19}$$

Suppose Victor first performs a von Neumann measurement on particle 1 and then on 3 and in both cases let the outcomes be $|y\rangle_1$ and $|y\rangle_3$. He can send the classical information (one cbit) to Alice and (one cbit) to Bob who can in turn find their particles in the original state exactly or up to a rotation operator, respectively. Thus the final state after von Neumann measurements is given by

$$\begin{aligned}
&|y\rangle_1\langle y|y\rangle_3\langle y|\Psi^+\rangle_{34}\langle\Psi^+|\Psi^-\rangle_{12}\langle\Psi^+|\Psi\rangle_{12345} = \\
&-\frac{1}{4}|y\rangle_1 \otimes |\Psi\rangle_2 \otimes |y\rangle_3 \otimes (\sigma_z)|\Psi\rangle_4 \otimes |\Psi\rangle_5.
\end{aligned} \tag{20}$$

Since particles 2, 4, and 5 are in possession of Alice, Bob and Carla, respectively it is clear from (20) that each of them acquire a perfect copy of the unknown state. On the other hand if Victor's outcome for particle 1 and 3 are $|x\rangle_1$ and $|x\rangle_3$, then the resulting state after communicating one bit to Alice and one to Bob is given by

$$\begin{aligned}
&|x\rangle_1\langle x|x\rangle_3\langle x|\Psi^+\rangle_{34}\langle\Psi^+|\Psi^-\rangle_{12}\langle\Psi^+|\Psi\rangle_{12345} = \\
&-\frac{1}{4}|x\rangle_1 \otimes |\Psi_\perp\rangle_2 \otimes |x\rangle_3 \otimes (-\sigma_z)|\Psi_\perp\rangle_4 \otimes |\Psi\rangle_5.
\end{aligned} \tag{21}$$

In this case Alice gets a complement copy, Bob gets a complement copy (up to a rotation operator) and Carla gets the original state. Similarly, depending on the outcomes of the measurement process on particle 1 and 3 two other possibilities exist. In general when Victor finds both particles in the basis $|y\rangle$, then we have 2 copies with probability 1/16 and when both particles are found in the basis $|x\rangle$, we have 2 complement copies with probability 1/16. However, if Victor finds particle 1 in the basis $|x\rangle$ and 3 in $|y\rangle$ (or vice versa) then we have a copy and a complement copy (all are up to doing nothing or a rotation operation) with probability 1/8. Thus the above protocol is able to produce two perfect copies or two complement copies or a copy and a complement copy (including original one) of an unknown state with the help of Victor. Since our protocol would work for all Bell-state outcomes, therefore, the probability of producing 2 clones (upto a rotation operator) is 25% and of producing 2 complement copies is 25% and 1 clone and 1 complement is 50%. It is a curious interplay between dual quantum and classical channel which is surprisingly enough able to produce *one to two* and *one to three* perfect copies and complement copies. The classical information (1 cbit) send by Victor is not enough to create a perfect copy from some arbitrary state *unless they have shared one half of the particle from the entangled state*. Here, again left-over entanglement after the teleportation process plays an important role in sending infinite bits from a state preparer. The present results can be generalised to produce multiple copies and complement copies using multiparticle entanglement source and multiparties. One can ask what is the useful resource to produce N copies or N complement copies (plus 1 original). It can be argued that we need $n = 2N$ particle entanglement source and $N + 1$ parties to be involved. The entangled source has to be distributed in such a way that the first and the last person possess one particle each and all intermediate parties possess 2 particles from the entangled source.

To conclude this paper, we have proposed a protocol which can produce perfect copies and orthogonal-complement copies of an unknown state with the help of dual quantum and classical channel (including minimal communication from state preparer). Though, in principle the state preparer knows what the state is he never reveals the information about the unknown state. This protocol uses the minimal amount of additional information (1 cbit) to produce a copy or a complement copy. This is the first time we are able to produce perfect copies and complement copies of an unknown state within quantum theory (which takes into account measurement processes). It is an open problem in quantum information theory how can one produce a copy or a complement copy of an unknown state within a quantum computer, where it is highly desirable to copy the information again and again. At least, we have shown that it is possible to do so perfectly with a minimal assistance from the state preparer. Our protocol provides an insight on what extra resource do we need to achieve perfect cloning and complementing. We hope this will be a practical way of copying and complementing quantum states in future. Also, the present work could have some application in quantum communication complexity [26] and in quantum bit commitment protocols [27].

I wish to thank S. Braunstein useful discussions. I thank V. Bužek and A. Chefles for going through my paper and sending their comments at some stage or the other. I thank L. M. Duan for useful discussions concerning the “no-cloning” theorem and teleportation process. I wish to thank D. Dieks for encouragements. Also, I thank P. Kok for a careful reading of the manuscript. I gratefully acknowledge the financial support from European Physical Science Research council (EPSRC).

-
- [1] W. K. Wootters and W. H. Zurek, *Nature (London)* **299**, 802 (1982).
 - [2] D. Dieks, *Phys. Lett. A* **92**, 271 (1982).
 - [3] H. P. Yuen, *Phys. Lett. A* **113**, 405 (1986).
 - [4] G. M. D’Ariano and H. P. Yuen, *Phys. Rev. Lett.* **76**, 2832 (1996).
 - [5] H. Barnum, C. M. Caves, C. A. Fuchs, R. Jozsa, and B. Schumacher, *Phys. Rev. Lett.* **76**, 2818 (1996).
 - [6] V. Bužek and M. H. Hillery, *Phys. Rev. A* **54**, 1844 (1996).
 - [7] M. H. Hillery and V. Bužek, *Phys. Rev. A* **56**, 1212 (1997).
 - [8] V. Bužek, S. L. Braunstein, M. Hillery and D. Bruß, *Phys. Rev. A* **56**, 3446 (1997).
 - [9] D. Bruß, D. P. Vincenzo, A. K. Ekert, C. A. Fuchs, C. Macchiavello and J. Smolin, *Phys. Rev. A* **57**, 2368 (1998).
 - [10] L. M. Duan and G. C. Guo, *Phys. Rev. Lett.* **80**, 4999 (1998).
 - [11] N. Gisin and S. Massar, *Phys. Rev. Lett.* **79**, 2153 (1997).
 - [12] N. Cerf, *Asymmetric quantum Cloning machines in any dimension*, quant-ph/9805024; quant-ph/9804011.
 - [13] A. Chefles and S. M. Barnett, *J. Phys. A* **31**, 10097 (1998).
 - [14] A. Chefles and S. M. Barnett, *Strategies and Network for state-Dependent Quantum Cloning*, quant-ph/9812035.
 - [15] A. K. Pati, *Multiple quantum clones can not be superposed*, Preprint (1999).
 - [16] A. K. Pati, *Quantum superposition of multiple clones and the novel cloning machine*, quant-ph/9903038.
 - [17] A. K. Pati, *New limitations in quantum information processing*, Preprint (1999).
 - [18] V. Bužek, M. Hillery and R. F. Werner, LANL Report, quantum-ph/9901053.
 - [19] N. Gisin and S. Popescu, LANL Report, quant-ph/9901072.
 - [20] R. Jozsa, *Geometric Issues in Foundations of Science*, Eds. S. Huggett et al, Oxford Univ. Press, 1997.
 - [21] C. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
 - [22] D. Bouwmeester, J. W. Pan, K. Mattle, M. Eible, H. Weinfurter, and A. Zeilinger, *Nature* **390**, 575 (1997).
 - [23] V. Braginsky, Y. I. Vorontsov, and K. S. Thorne, *Science* **209**, 547 (1980).
 - [24] D. M. Greenberger, M. A. Horne and A. Zeilinger, *Bell’s Theorem, Quantum Theory and Conceptions of the Universe* Edt. M. Kafatos, (Kluwer Academic Publishers, 1989, The Netherlands)
 - [25] A. Karlsson and M. Bourennane, *Phys. Rev. A* **58**, 4394 (1998).
 - [26] E. Kushilevitz and N. Nisan, *Communication Complexity*, Cambridge University Press, 1997.
 - [27] H. K. Lo and H. F. Chau, *Phys. Rev. Lett.* **78**, 3410 (1997); D. Mayers, *Phys. Rev. Lett.* **78**, 3414 (1997).